

Chronic subdural hematoma

The essence of **chronic subdural hematoma** (CSDH) is bleeding from the bridging veins in patients with variously expressed cortical **atrophy** (elderly patients, alcoholics), as well as in infants. CSDH was first described by Rudolf Virchow in 1857 as "*pachymeningitis haemorrhagica interna*" [1], after which Wilfred Totter described the basics of CSDH pathophysiology and thus defined that hemorrhage from the bridging veins is its root cause and named it "*subdural hemorrhagic*" [2].

It is important to adequately distinguish CSDH from acute subdural hematoma. Compared with CSDH, acute SDH typically occurs in younger individuals primarily as a result of more severe cranio traumas and is associated with symptomatic compression of the brain parenchyma, which typically manifests within 72 hours. Conversely, in the case of CSDH, bleeding is **clinically asymptomatic for a longer period of time**, on the order of several weeks or months, most often occurring after **minor injuries**. The incidence of CSDH in the population is around 1.72 / 100,000 inhabitants per year, in the population aged 70–79 this value increases to 7.35 / 100,000 per year, thus further confirming the fact of a higher risk of developing CSDH in the elderly.

Risk factors

The basic risk factors include **cerebral atrophy**, associated with older age. In direct proportion to aging, the space between the calva and the parenchyma itself increases from the original 6% to 11%. This allows the brain to move sideways, so in the event of trauma, bridging veins can be more easily damaged, resulting in CSDH. [3]. However, trauma alone is not the only cause of CSDH, patients with **anticoagulant medication**, **alcoholism** or **epilepsy** are also affected (up to 10% of patients with CSDH are alcoholics or people with a history of seizures). [4].



Subdural hematoma

Pathophysiology and symptomatology


Damage to the bridging veins (eg in the case of trauma) results in bleeding into the subdural space, with a thin layer of fibrin and fibroblast forming around the hematoma surrounding the hematoma. Fibrin and fibroblasts further proliferate and migrate, forming a kind of membrane (approximately 4 days after the incident), which then progressively enlarges. Some CSDH is **resorbed spontaneously**, but most grows in volume and results in **clinically symptomatic CSDH** [5], neurologically manifested by headache, confusion or, at a more advanced stage, contralateral hemiparesis.

Diagnosis

CT is used to image the hematoma. It clearly displays the resulting hematoma. It is typically a hypodense extracerebral bed, see picture.

Therapy

In the case of large CSDHs without expansive tendencies and clinically good patient status, it is not necessary to evacuate the hematoma, **conservative treatment** consists in close monitoring of the patient [6]. In the case of a progressive symptomatic lesion, we indicate a **trepanation return**, ie the evacuation of coagulated blood and the introduction of temporary subdural drainage from the return site under local anesthesia. In some cases, **craniotomy** is used, especially in patients with recurrent hematomas, resp. in persons showing blood re-accumulation after trepanation return, or the presence of a solid hematoma. Rehydration of the patient and care for the patency of the drainage under strictly sterile conditions is important, while a control CT is performed after 3 days and the drain is removed if the finding is satisfactory. [7][8]



Chronic subdural hematoma in an elderly patient with evident cortical atrophy. The blue arrow points to the hematoma site, the red one points to a trepanning hole.

Links

Related articles

- Craniocerebral trauma

- Acute subdural hematoma
- Subdural hematoma
- Epidural hematoma
- Subarachnoid hemorrhage

References

- SAMEŠ, M, et al. *Neurochirurgie*. 1. edition. Praha : Jessenius Maxdorf, 2005. ISBN 80-7345-072-0.

Reference

- ws: Chronický subdurální hematom
1. ADHIYAMAN, V. Chronic subdural haematoma in the elderly. *Postgraduate Medical Journal*. 2002, y. 916, vol. 78, p. 71-75, ISSN 0032-5473. DOI: 10.1136/pmj.78.916.71 (<http://dx.doi.org/10.1136%2Fpmj.78.916.71>).
 2. FURLOW, LEONARD T.. CHRONIC SUBDURAL HEMATOMA. *Archives of Surgery*. 1936, y. 4, vol. 32, p. 688, ISSN 0004-0010. DOI: 10.1001/archsurg.1936.01180220114007 (<http://dx.doi.org/10.1001%2Farchsurg.1936.01180220114007>).
 3. ELLIS, G L. Subdural hematoma in the elderly. *Emerg Med Clin North Am [online]*. 1990, vol. 8, no. 2, p. 281-94, Available from <<https://www.ncbi.nlm.nih.gov/pubmed/2187683>>. ISSN 0733-8627.
 4. JONES, S. A prospective study of chronic subdural haematomas in elderly patients. *Age and Ageing*. 1999, y. 6, vol. 28, p. 519-521, ISSN 1468-2834. DOI: 10.1093/ageing/28.6.519 (<http://dx.doi.org/10.1093%2Fageing%2F28.6.519>).
 5. MUNRO, DONALD. SURGICAL PATHOLOGY OF SUBDURAL HEMATOMA. *Archives of Neurology & Psychiatry*. 1936, y. 1, vol. 35, p. 64, ISSN 0096-6754. DOI: 10.1001/archneurpsyc.1936.02260010074005 (<http://dx.doi.org/10.1001%2Farchneurpsyc.1936.02260010074005>).
 6. NAGANUMA, Hirofumi – FUKAMACHI, Akira – KAWAKAMI, Motomasa. Spontaneous Resolution of Chronic Subdural Hematomas. *Neurosurgery*. 1986, y. 5, vol. 19, p. 794-798, ISSN 0148-396X. DOI: 10.1227/00006123-198611000-00013 (<http://dx.doi.org/10.1227%2F00006123-198611000-00013>).
 7. SMELY, C. – MADLINGER, A. – SCHEREMET, R.. Chronic subdural haematoma ? a comparison of two different treatment modalities. *Acta Neurochirurgica*. 1997, y. 9, vol. 139, p. 818-826, ISSN 0001-6268. DOI: 10.1007/bf01411399 (<http://dx.doi.org/10.1007%2Fbf01411399>).
 8. BENDER, M. B. – CHRISTOFF, N.. Nonsurgical Treatment of Subdural Hematomas. *Archives of Neurology*. 1974, y. 2, vol. 31, p. 73-79, ISSN 0003-9942. DOI: 10.1001/archneur.1974.00490380021001 (<http://dx.doi.org/10.1001%2Farchneur.1974.00490380021001>).