

Hemoglobin and its derivatives (1. LF UK, VL, Practical exercises in biochemistry)

This article is intended for teaching Practical Exercises in Biochemistry at the 1st Faculty of Medicine, Charles University. There is a basic article in WikiScripts on the Hemoglobin page .

Hemoglobin is a red blood pigment that transports oxygen from the lungs to the tissues and transports CO₂ and protons from peripheral tissues to the respiratory system.

The hemoglobin concentration in a healthy adult male is approximately 150 g / l, in an adult female about 140 g / l. One gram of hemoglobin can bind up to 1.34 ml of oxygen.

Content

- 1Hemoglobin structure
- 2Hemoglobin in the blood
- 3Causes of anemia
- 4Principle of hemoglobin determination in blood
- 5Hemoglobin in urine
 - 5.1Determination of hemoglobin in urine
- 6Hemoglobin in stool - occult bleeding
- 7Hemoglobin derivatives
 - 7.1Oxyhemoglobin and deoxyhemoglobin
 - 7.2Carbaminohemoglobin
 - 7.3Methemoglobin
 - 7.4Carbonyl hemoglobin
 - 7.5Spectrophotometry of hemoglobin derivatives
- 8Iron
 - 8.1Iron metabolism
 - 8.2Examination of iron metabolism
 - 8.2.1Determination of iron in serum
 - 8.2.2Serum transferrin and iron binding capacity
 - 8.2.2.1Transferrin saturation
 - 8.2.3Ferritin and hemosiderin
 - 8.2.4Transferrin receptor
- 9Disorders of iron metabolism
 - 9.1Iron deficiency (sideropenia)
 - 9.2Excess iron
 - 9.3Iron poisoning
- 10Links
 - 10.1Reference
 - 10.2References

Hemoglobin structure

Hemoglobin It is a tetrameric protein formed by four subunits. The two and two subunits are always identical. There are four types of polypeptide chains, physiologically occurring hemoglobin, α , β , γ , and δ , which differ in the number and sequence of amino acids . The tetramer consists of two α chains and two other types of chains that indicate the character of the whole hemoglobin molecule. In adults, hemoglobin A predominates, with two β chains (146 amino acids) involved in addition to two α chains (141 amino acids).

Each subunit includes a polypeptide chain to which one heme is covalently attached . The basis of the heme molecule is protoporphyrin, formed by 4 pyrrole nuclei connected by methenyl bridges with centrally bound iron. Heme iron is a total of six bonds - it is connected to the nitrogen atoms of the pyrrole nuclei by four coordination bonds. By another coordination valence, iron binds to the imidazole group of the amino acid histidine in the globin chain. The sixth valence Fe is determined for the molecule of oxygen (O₂).

Hemoglobin in the blood

[ edit embedded article]

Determination of hemoglobin in the blood is one of the most basic laboratory tests. Blood hemoglobin is the main criterion for assessing whether it is anemia . The term anemia is used when the hemoglobin or erythrocytes fall below the lower limit of physiological levels. Anemia is a very common clinical finding. This is a condition that leads to a reduction in oxygen binding capacity and a consequent tissue respiratory disorder.

Causes of anemia

Anemia occurs when erythropoiesis is unable to meet the requirements for new red blood cells. It develops as a result of blood loss or increased loss of red blood cells or insufficient red blood cell production. The following is a list of some specific causes of anemia:

1. *Anemia from increased blood loss* :
 - Acute blood loss.
 - Chronic blood loss.
2. *Anemia due to increased erythrocyte breakdown* (hemolytic conditions).
 - Autoimmune hemolytic anemia (presence of antibodies against own erythrocytes).
 - Erythrocyte membrane disorder (deviation in erythrocyte membrane composition).
 - Hereditary erythrocyte enzyme defects (pyruvate kinase, glucose-6-phosphate dehydrogenase).
 - Unstable hemoglobin - hemoglobinopathies (eg hemoglobin S in sickle cell disease).
3. *Anemia from decreased erythrocyte production* .
 - Lack of substances needed for erythropoiesis (iron deficiency, vitamin B12 deficiency, folic acid deficiency, erythropoietin deficiency - chronic renal diseases, lack of other substances such as vitamins B1, B6).
 - Anemia due to chemical, physical and radiation damage.
 - Anemia in chronic inflammatory, infectious and cancerous diseases.

Elevated hemoglobin levels may be a sign of dehydration or chronic decreased pulmonary ventilation. Rarely, it can be caused by some myeloproliferative conditions, such as polycythemia vera .

Principle of hemoglobin determination in blood

Oxidation of hemoglobin to methemoglobin:

HbFe II	+	[Fe III (CN) 6] 3–	→	HbFe III	+	[Fe II (CN) 6] 4–
Hemoglobin				Methemoglobin		

Conversion of methemoglobin to cyanomethemoglobin:

HbFe III	+	CN -	→	HbFe III CN	
Methemoglobin				Cyanmethemoglobin	

The photometric determination is based on the oxidation of ferrous iron in hemoglobin with potassium ferrocyanide to ferric iron. The resulting methemoglobin is converted to a very stable cyanomethemoglobin in a further reaction with potassium cyanide with a single broad absorption maximum in the visible region at 540 nm.

Assessment: The reference range for hemoglobin in the blood (B hemoglobin) for an adult male is 130-180 g / l and for a female 120-160 g / l.

Hemoglobin in urine

[ edit embedded article]

Up to a million erythrocytes per day are excreted in the urine of completely healthy people. This very small amount cannot be demonstrated by conventional chemical tests. Occurrence of a larger number of erythrocytes (hematuria , erythrocyturia) or penetration of free hemoglobin, or muscle myoglobin, into definitive urine (hemoglobinuria or myoglobinuria) is almost always a pathological finding. We observe macroscopic hematuria with the naked eye; the urine is pinkish (comparable to water from washed meat) and hemoglobin can be detected spectroscopically in it. There is at least 1 g of hemoglobin per liter in the urine. In massive hemoglobinuria, the urine may be dark beer (degradation of hemoglobin to hematin). Microscopic hematuriait can only be detected biochemically.

Determination of hemoglobin in urine

[ edit embedded article]

Hemoglobin catalyzes, like peroxidase , the oxidation (dehydrogenation) of some substrates (eg benzidine derivatives) by hydrogen peroxide:




However, it is not an enzyme activity (catalysis is conditioned by heme iron) and therefore it is not lost even after heat denaturation. We are talking about pseudoperoxidase activity , which is used for sensitive but non-specific evidence of hemoglobin or trace amounts of blood. It is preferable to use a chromogenic substrate to monitor the reaction, i.e., a substance that provides a markedly colored product by dehydrogenation (often benzidine or its non-carcinogenic derivatives, aminophenazone, etc.).

The reagent zone of the diagnostic strips contains a chromogen (eg tetramethylbenzidine) with stabilized hydrogen peroxide (eg cumene hydroperoxide). In the presence of free hemoglobin (hemoglobinuria), the indication zone turns uniformly blue. If erythrocytes (erythrocyturia) are present in the urine , intensely green-blue dots to spots form.

Hemoglobinuria can be encountered in intravascular hemolysis . Damage to the glomerular membrane (glomerular hematuria) and bleeding from any part of the urinary tract lead to more frequent erythrocyturia . It is often found in urinary tract infections , urolithiasis and urogenital tract tumors .

In addition to hemoglobin , myoglobin also provides a pseudoperoxidase response , which can be excreted in the urine during skeletal muscle breakdown (rhabdomyolysis , crush syndrome). The positivity of the test may also be due to peroxidases of leukocytes or certain bacteria, yeasts or fungi, which may occur in the urine, especially in urinary tract infections . To rule out the possibility of a false positive reaction due to cellular peroxidases, the reaction must be performed with boiled urine.



Contamination of the sampling vessel with strong oxidizing agents also causes a false positive reaction. On the other hand, the presence of strong reducing substances (eg ascorbic acid) can slow down or even stop the pseudoperoxidase reaction and thus cause false negative results.

Hemoglobin in stool - occult bleeding

Occult (hidden) bleeding (OK) is a term for chemical detection of blood in stool that has a normal macroscopic appearance. Demonstration of occult stool bleeding is used to detect the early stages of colorectal cancer , when radical and effective treatment is still possible. It also allows the detection of precancerous diseases of the colon, especially adenoma polyps.

In principle, two methodological approaches are used to capture occult bleeding:

- Methods based on the demonstration of pseudoperoxidase activity of hemoglobin. The principle is the same as for the determination of hemoglobin in urine . The patient must maintain a diet for 3 days before the examination, exclude uncooked meat, salami, bananas, tomatoes from the diet, and must not take drugs containing ascorbic acid or acetylsalicylic acid. The patient then takes samples from three consecutive stools and applies them to chromogen-saturated test cards. The evaluation is performed in the office or in the laboratory by adding a solution of hydrogen peroxide and subtracting the color.

See the Haemoccult page for more information .

- Other methods are based on the immunochemical detection of hemoglobin with an anti -human hemoglobin antibody. Immunochemical methods are more sensitive and specific, there is no need to follow a diet before the examination.

See the Stool Occult Bleeding Tests page for more information .

Positive results must be verified by other diagnostic methods.


Hemoglobin derivatives

[[edit embedded article](#)]

Hemoglobin derivatives include the following types:

Oxyhemoglobin and deoxyhemoglobin

[[edit embedded article](#)] Oxyhemoglobin and deoxyhemoglobin Oxygen - carrying hemoglobin is referred to as oxyhemoglobin (oxyHb) . Each Hb molecule can bind 4 molecules of oxygen. After the release of oxygen, we speak of deoxyhemoglobin (deoxyHb) . In both forms, iron is divalent because only Fe II + -containing hemoglobin can reversibly bind and transport the oxygen molecule. Oxygenation of the hemoglobin molecule changes the electronic state of the Fe II + -hem complex, which results in a change in the color of the dark red hue typical of venous blood to the bright red color of arterial blood. In the human body, about 98.5% of oxygen is bound to hemoglobin.



Carbaminohemoglobin

[[edit embedded article](#)]

Carbaminohemoglobin is a hemoglobin to which CO₂ is bound . Carbon dioxide binds to the globin chain of hemoglobin . The binding of CO₂ to hemoglobin reduces the affinity of hemoglobin for oxygen.

Methemoglobin

[[edit embedded article](#)]

Methemoglobin (metHb; also *hemoglobin* or *ferrihemoglobin*) is characterized by the presence of ferric iron , which is formed by the oxidation of ferrous iron in hemoglobin . Methemoglobin loses its ability to reversibly bind oxygen . In its place, Fe III + binds a water molecule through the sixth coordination bond. The color of methemoglobin is


chocolate brown.

Methemoglobin is also present physiologically in small amounts in erythrocytes (about 1–3% of the total hemoglobin concentration). This is mainly due to the effect of nitrite, which is formed from nitrates contained in food. The reverse reduction of methaemoglobin to hemoglobin is mainly ensured by NADH-dependent cytochrome-b 5 reductase (also methemoglobin reductase). A minor role is played by NADPH-dependent methemoglobin reductase, which is dependent on the supply of NADPH from the pentose cycle and on the presence of another electron transporter (eg flavin). Non-enzymatic mechanisms include the action of glutathione and ascorbic acid . Methemoglobinemia - origin and therapy Elevated blood levels of methaemoglobin are called methaemoglobinaemia . The causes are different:

- Hereditary methaemoglobinaemia is usually caused by a congenital defect in NADH-dependent methaemoglobin reductase or the presence of abnormal hemoglobin M.
- Acquired methaemoglobinaemia is the most common form of methaemoglobinaemia. May be caused by oxidizing agents :
 - poisoning by certain substances (nitrobenzene , aniline and its derivatives - eg some dyes),
 - by the action of some drugs (local anesthetics - benzocaine , then phenacetin , sulfonamides),
 - increased content of nitrates and nitrites in water and food.

Newborns are particularly sensitive to the increased content of these substances due to the immaturity of the reduction systems and the increased proportion of fetal hemoglobin, which is more easily oxidized. Methaemoglobinaemia is manifested by cyanosis with a characteristic gray-brown tint and hypoxia.

1. vznik methemoglobinu



2. redukce methemoglobinu na hemoglobin



Symptoms of methemoglobinemia

Methemoglobin values	Symptoms
0-2%	normal value
<10%	cyanosis
<35%	cyanosis and other symptoms (headache, shortness of breath)
70%	lethal concentration

Part of the therapy of acquired methaemoglobinaemia is the administration of some reducing agents - methylene blue or ascorbic acid.

Carbonyl hemoglobin

[[🔗 edit embedded article](#)]

Carbonylhemoglobin (COHb, *carboxyhemoglobin*) is formed by the binding of carbon monoxide to hemoglobin . The bond formed is 250-300 times stronger than the oxygen bond . Carbonyl hemoglobin cannot transport oxygen and cellular hypoxia develops due to the blood 's reduced ability to carry oxygen . In excess oxygen, the binding of carbon monoxide to hemoglobin is reversible. Therefore, inhalation of O₂ is the most important in carbon monoxide poisoning .

COHb can also be found in small amounts in healthy people. For urban dwellers, values of around 2% are evident; for heavy smokers , COHb can rise to as much as 10% of total hemoglobin. Staying in an environment containing 0.1% CO for several minutes can increase the carbonyl hemoglobin concentration to 50%.

Carbon monoxide is formed during the imperfect combustion of fuels, it is also contained in the exhaust gases and in smoke during fires in closed rooms. Hemoglobin saturation curve

Symptoms of carbon monoxide poisoning

COHb values in%	Symptoms
10	shortness of breath on exertion
20-40	headache , shortness of breath, fatigue, vomiting
40-60	hyperventilation, tachycardia, syncope, convulsions
60-80	coma, death


Carbonyl hemoglobin is crimson red; even people with severe carbon monoxide poisoning tend to have "healthy" pink skin. Compared to hemoglobin, carbonyl hemoglobin is more resistant to chemical influences, it changes more slowly due to the action of various agents.

Spectrophotometry of hemoglobin derivatives

[[edit embedded article](#)] Absorption spectra of hemoglobin and its derivatives Hemoglobin and its derivatives have characteristic absorption spectra in the visible region of light, which are used for their analysis and rapid identification. Significant absorption maxima in the range of 400–430 nm, the so-called Soret band, are typical for all hemoproteins. Other absorption peaks are significantly lower. Oxyhemoglobin is characterized by two incompletely separated maxima in the region of 540 and 578 nm. Deoxyhemoglobin has a single absorption maximum at 555 nm. The main absorption maximum of methaemoglobin is at 630 nm and the second faint peak at 500 nm is pH dependent. By reacting methemoglobin with potassium cyanide, the maximum disappears at 630 nm, as cyanomethemoglobin is formed. The decrease in absorbance at 630 nm is proportional to the methaemoglobin concentration. Cyanmethemoglobin shows a broad absorption maximum at 540 nm, which is used to determine blood hemoglobin concentration. The carbonyl hemoglobin spectrum is similar to that of oxyhemoglobin, but the position of the peaks is shifted toward lower wavelengths.

Absorption maxima of hemoglobin and its derivatives

Hemoglobin derivative	Absorption maxima [nm]
Hemoglobin reduced	431, 555
Oxyhemoglobin	414, 540, 578
Methemoglobin	404, 500, 630
Carbonyl hemoglobin	420, 538-540, 568-569
Cyanmethemoglobin	421, 540



Determination of carbonyl hemoglobin: Determination of carbonyl hemoglobin spectrophotometry Determination of carbonyl hemoglobin in the blood is one of the basic toxicological examinations. It is an objective criterion in the assessment of acute and chronic carbon monoxide poisoning.

- *Spectrophotometric evaluation* . Carbonyl hemoglobin can be determined rapidly spectrophotometrically by subtracting the absorption maximum shift of the diluted blood from 586 nm . The shift of the maximum in the spectrum depends on the ratio of COHb and O₂ Hb in the sample.
- *Reactions with tannin* . As a guide, carbonyl hemoglobin can be determined by reaction with tannin or Ajatin (from about 10% COHb). Tannin forms a strawberry red precipitate in the presence of carbonyl hemoglobin. In the absence of carbonyl hemoglobin, the color of the precipitate is brownish gray.
- *Acid-base balance analyzers* . The analysis of the most toxicologically important hemoglobin derivatives COHb and metHb is also enabled by modern acid-base balance analyzers, which have a built-in photometric system for their measurement.

Iron

[[edit embedded article](#)]

Iron is one of the most important elements in the human body. The adult body contains more than 70 mmol (4.0-4.5 g) of iron. In women, this amount is lower than in men, which is attributed to blood loss during menses .

Distribution of iron in the body

Form	Function	Protein	Amount in g
Active iron	oxygen transport	hemoglobin	2.5-3.0
		myoglobin	0.3
	electron transfer	cytochromes , cytochrome oxidase	0.2
	decomposition of hydrogen peroxide	catalase , peroxidase	
Storage iron		ferritin , hemosiderin	0.8-1.0
Transport iron		transferrin	0.003

Iron metabolism

The presence of iron is essential for cell function. As a part of heme it participates in the transport of oxygen and as a part of cytochromes it conditions the transfer of electrons in the respiratory chain. An undesirable effect of iron as a transient and highly reactive element is its participation in radical reactions, in which so-called reactive oxygen species are formed . These can damage cell membranes, proteins and DNA.

Iron is absorbed as Fe²⁺ by active transport in the duodenum and upper jejunum in two ways:

1. porphyrin-bound Fe in the form of a stable lipophilic complex;
2. Fe^{II} + - water soluble chelates.

Only a small part is absorbed in ionized form.

The diet averages 10-50 mg of iron per day, but only 10-15% is absorbed. Heme (meat) compounds absorb better, non-heme Fe in the plant diet much worse. In addition, plants contain oxalates, phytates, tannins and other phenolic compounds which form insoluble or chelated complexes with Fe which are difficult to absorb. Ascorbic acid, on the other hand, improves iron absorption.

After uptake by the intestinal mucosa, part of the iron is incorporated into the storage form - ferritin in intestinal cells. Part of the absorbed iron passes into the plasma, where it is transported in binding to transferrin. The protein ferroportin (it is also found in the membrane of macrophages and hepatocytes) plays an important role in the transfer of iron across the basolateral membrane of enterocytes. It is the main site of regulation of iron homeostasis in the body. A key regulatory factor is the hepcidin protein, which is synthesized in the liver. By binding to ferroportin, it inhibits the transport of iron from cells and thus contributes to its sequestration in them. Hepcidin levels increase with inflammation. Hepcidin is also partly responsible for anemia of chronic diseases. Mutations in the hepcidin gene lead to juvenile hemochromatosis type 2B.


Plasma iron is captured by target tissue cells via the transferrin receptor and is either incorporated into heme or stored in ferritin. The use of the specific transport protein transferrin and the ferritin storage protein for iron storage represents protective mechanisms to prevent the toxic effects of redox active iron.

During desquamation of dead mucosal cells, unused iron leaves the stool together with unabsorbed iron.

Examination of iron metabolism

In practice, we commonly encounter diseases associated with changes in iron metabolism and utilization. Laboratory tests of iron metabolism include the following tests:

- iron in serum
- serum transferrin and iron binding capacity
- serum ferritin
- transferrin receptor



These parameters are important diagnostic indicators for demonstrating a decrease or increase in iron stores even in stages that are not accompanied by significant clinical manifestations.

Determination of iron in serum

Principle of determination of iron in serum Colorimetric methods, atomic absorption spectrophotometry and other special techniques are used to determine serum iron. The most used are photometric methods based on the reaction of iron with a complexing agent. All procedures include the following steps: Ferrozine

1. Release of Fe³⁺ from transferrin binding by acids or surfactants (eg HCl).
2. Reduction of Fe³⁺ to Fe²⁺, which is necessary for the reaction with the complexing agent. Ascorbic acid, for example, is used for the reduction.
3. Reaction of Fe²⁺ with a complexing agent containing reactive groups -N = C - C = N- to form a color complex. Metal ions form chelates with two nitrogen atoms. Currently, mainly two complexing agents are used - bathofenanthroline and ferrozine (3- (2-pyridyl) -5,6-bis (4-sulfophenyl) -1,2,4-triazine - PST, protected name FerroZine®), which has higher absorption coefficient and is more soluble in water.

Evaluation

Serum iron concentrations are subject to circadian rhythm and are influenced by other factors. This limits the diagnostic significance of this parameter. It is a poor indicator of tissue iron stores and should always be considered in combination with serum transferrin and iron binding capacity. Decreased concentrations are accompanied by iron deficiency, caused for example by large or repeated blood loss, insufficient dietary iron intake or impaired absorption. The finding is not specific, as reduced levels are also encountered in acute infection or chronic inflammatory diseases (iron transfer to tissues). High iron levels occur in hemochromatosis (see below), in iron overdose or intoxication, in increased erythrocyte breakdown, and in some liver diseases.

Reference values

men: 9-29 μmol / l
women: 7-28 μmol / l

Serum transferrin and iron binding capacity

Iron is transported by the blood in binding to a specific protein with β 1 -electrophoretic mobility - transferrin, which is synthesized in the liver. The rate of its formation is inversely proportional to the body's iron stores; it increases with iron deficiency and decreases with excess. The biological function of transferrin is the ability to easily form non-toxic iron complexes and transfer Fe absorbed by the small intestinal mucosa to the bone marrow or storage forms (ferritin or hemosiderin). Each transferrin molecule binds two Fe³⁺ atoms (1 g transferrin binds 25.2 μmol iron). Transferrin can be determined directly by immunochemical methods or indirectly as the ability of transferrin to bind iron - the so-called iron binding capacity. Total *iron binding capacity* (TIBC) is the amount of iron that transferrin is able to bind when all binding sites are occupied. Usually only 1/3 of transferrin-bound capacity is saturated with iron. Free transferrin without bound iron represents the free binding capacity (2/3 of transferrin) that is available for iron transport in increased demands.

Conversion between transferrin concentration and total binding capacity:

Total binding capacity [$\mu\text{mol} / \text{l}$] = transferrin [g / l] \cdot 25.2 .

The reference range for serum transferrin concentration (S-transferrin) is 2.0–3.6 g / l and for a total binding capacity is 50–70 $\mu\text{mol} / \text{l}$.

Transferrin saturation

From the values of iron and transferrin concentration, we can calculate transferrin saturation (TfS) , which is defined as the ratio of serum iron concentration to total iron transfer capacity for transferrin. This is a sensitive parameter for detecting latent iron deficiency.

Extra open brace or missing close brace

Evaluation of transferrin saturation

physiological values: 25-50%

iron deficiency saturation reduction: <15%

increase in saturation with excess iron:> 50%

Ferritin and hemosiderin

Ferritin is the most important storage protein for iron. The ferritin molecule is adapted to bind large amounts of Fe^{3+} in a soluble and non-toxic form to the body. Ferritin consists of an outer protein shell of 24 subunits - apoferritin (Mr 440,000), delimiting a cavity in which up to 4500 iron atoms can be concentrated in the form of ferric oxyhydroxide ($\text{FeO} \cdot \text{OH}$) in microcrystalline form with phosphates ($\text{FeO} \cdot \text{OPO}_3\text{H}_2$). The entry and exit of iron atoms is enabled by the pores between the individual subunits of the ferritin shell. Normally, its capacity is used by about 20%. It is stored in cells in the liver, spleen and intestinal mucosa.

Ferritin is found in very low concentrations in the blood serum. Serum ferritin concentrations are a measure of the body's iron stores. Low concentrations indicate depletion of the body's total iron reserve and serve to detect iron deficiency anemia early in the prelatent phase. Elevated ferritin concentrations are an accompanying phenomenon of high tissue iron stores. We also encounter them in many patients with liver disease, some malignancies (tumor marker) or inflammatory diseases (acute phase positive reactant).

The reference range for serum ferritin (S-ferritin) is 30-300 $\mu\text{g} / \text{l}$ for men and 20-120 $\mu\text{g} / \text{l}$ for women.

Hemosiderin is another storage protein for iron. It is formed by aggregation of denatured ferritin with other components. It forms particles with a size of 1 to 2 μm , which are visible in a light microscope when iron staining is used. Hemosiderin contains more iron than ferritin, but is difficult to obtain due to its poor water solubility. It is formed when the amount of iron in the body exceeds the storage capacity of ferritin.

Transferrin receptor

Iron transported by blood by transferrin is taken up by cells via a specific transferrin receptor (TfR). It is on the surface of all cells at some stage of development, but is most expressed on the surface of red cell precursors in the bone marrow. TfR is a transmembrane protein that is made up of two identical disulfide-linked subunits. By separating the extracellular domains of the receptor, the so-called soluble fraction of the transferrin receptor (sTfR) , which may be in the form of a dimer or monomer, is released into the circulation. Cells respond to reduced iron stores by synthesizing increased levels of transferrin receptors.


An increase in sTfR is a reliable indicator of iron deficiency for hematopoiesis . Elevated sTfR levels are found in *iron deficiency anemias* or *hemolytic anemias* . Determination of sTfR is valuable in anemic patients in whom ferritin is elevated due to an acute phase reaction. Determination of sTfR levels can also be used in patients with bone marrow transplantation to monitor the course of erythropoiesis.

Immunochemical methods are used for the determination.


Disorders of iron metabolism

[ edit embedded article]

Iron deficiency (sideropenia)



Ferritin structure diagram
Průchová Petr, Maršal, 2011



Transferrin receptor
Průchová Petr, Maršal, 2011

Iron deficiency in the body is usually caused by insufficient absorption from the intestine or chronic blood loss. It can result in sideropenic anemia (hypochromic microcytic anemia) , which is one of the most common hematological diseases . However, anemia is usually a late symptom of progressive sideropenia. It does not appear in the blood picture until the iron has almost completely disappeared. Therefore, it is necessary to detect iron deficiency at an early stage, which is not yet accompanied by anemia.

Based on the determination of the basic parameters of iron metabolism, we distinguish three degrees of deficiency:

- Prelatant iron deficiency is a sign of a state where there is a gradual decline in stocks, but the supply of iron to the bone marrow erythroblasts is not yet affected . In about half of the patients, serum ferritin levels are reduced below 12 µg / l.
- With latent iron deficiency , its reserves are essentially depleted. Ferritin is reduced below the lower limit of normal and is already accompanied at this stage by a reduction in serum iron levels and a reduced supply to bone marrow erythroblasts. The binding capacity for iron increases. A sensitive indicator of latent iron deficiency is a decrease in transferrin saturation below 15%. However, anemia is not developing yet.
- With manifest iron deficiency , anemia develops with hemoglobin levels falling below the lower limit of normal. Iron deficiency anemia is characterized by low serum iron and ferritin, and an increase in transferrin concentration (iron binding capacity). In contrast, in hemolytic anemias or in excess of iron, serum iron is increased, and at the same time the total iron binding capacity is reduced.

Laboratory finding in iron deficiency

Prelatant iron deficiency	Latent iron deficiency	Manifest iron deficiency
reduction of storage iron - decrease of ferritin	iron deficiency - decrease in ferritin	iron deficiency - decrease in ferritin
	reduction of serum iron	reduction of serum iron
	transferrin saturation drop below 15%	transferrin drop below 10%
	increasing the overall binding capacity for iron	increasing the overall binding capacity for iron
	increase in sTfR	increase in sTfR
		reduction in hemoglobin - anemia

Excess iron

The body is not equipped with an excretory pathway for iron, and therefore, under certain circumstances, excess iron may accumulate in the tissues. Early diagnosis can prevent tissue damage from excess iron. Iron overload usually develops very slowly. We distinguish 3 stages:

- In the stage of prelatent iron surplus , its content in the organs increases, but without exceeding their storage capacity.
- During the latent stage of iron overload , the storage capacity of the cells is exceeded, but the function of the organs is not yet impaired, the ferritin level and the serum iron level increase and the transferrin saturation rises above 55%.
- In the phase of manifest iron surplus , some organs are already damaged.

Laboratory finding of excess iron

Prelatent excess of iron	Latent excess iron	Manifest surplus of iron
increasing iron reserves - increasing ferritin	increasing iron reserves - increase in ferritin above 300 µg / l	increase in iron reserves - increase in ferritin (in case of severe impairment above 2000 µg / l)
	increase in serum iron	significant increase in serum iron
	increase in transferrin saturation above 55%	increase in transferrin saturation (may exceed 90% in severe disability)

Hemochromatosis

The accumulation of iron in the tissues is related to a disease we call hemochromatosis .

- Primary hemochromatosis is an inherited disease caused by increased resorption of iron in the gut. Excess iron is stored in parenchymal organs such as the liver, heart, pancreas, adrenal glands. It is toxic to the affected organs and disrupts their function by catalyzing chronic reactions leading to the formation of free radicals. The main clinical manifestations are skin hyperpigmentation, hepatosplenomegaly and diabetes mellitus .
- Secondary haemochromatosis may develop as a result of, for example, repeated transfusions, excessive iron intake or haemolytic anemia. In the biochemical picture we find increasing levels of ferritin and iron in the serum, the saturation of transferrin increases with its simultaneous decrease.

Iron poisoning

[ edit embedded article]

Children (tablets similar to lentils) are at risk of accidental ingestion of large quantities. The lethal dose for a child is 600 mg. For adults, iron intake of 40 mg / kg is toxicologically severe, intake of 60 mg / kg is fatal .

Symptoms include nausea, vomiting (including vomiting blood), abdominal pain, diarrhea (sometimes bloody). Large fluid losses cause shock, kidney failure and death. If the patient survives this phase of poisoning, he may become unconscious, convulsive and have failed after 12 hours. If he survives this second phase, the poisoning can have permanent consequences (intestinal damage).

Treatment of acute poisoning

- Gastric lavage.
- Use the nasogastral tube to deliver the chelating agent deferoxamine (5-10 g in 50-100 ml of water).
- Consider intravenous administration of desferoxamine to unbind the absorbed iron. A pinkish red complex of deferoxamine with iron appears in the urine. Treatment should be repeated until urine returns to normal .

Links

Reference

1. ↑ Jump up to: a b ŠVÍGLEROVÁ, Jitka. *Hemoglobin* [online]. Last revision 2009-02-18, [cit. 2010-11-11]. < <https://web.archive.org/web/20160416205421/http://wiki.lfp-studium.cz/index.php/Hemoglobin> >.
2. ↑ KITTNAR, Otomar and ET AL .. *Medical physiology*. 1st edition. Prague: Grada, 2011. 790 pp. 131. ISBN 978-80-247-3068-4 .
3. ↑ ŠVECOVÁ, D and D BÖHMER. Congenital and acquired methaemoglobinaemia and their treatment. *Journal of Czech doctors*. 1998, vol 137, pp. 168-170, ISSN 1803-6597.
4. ↑ RICHARD, Alyce M, James H DIAZ and Alan David KAYE. Reexamining the risks of drinking-water nitrates on public health. *Ochsner J* [online] . 2014, vol. 14, no. 3, pp. 392-8, also available from < <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4171798/?tool=pubmed> >. ISSN 1524-5012.
5. ↑ XU, F, KS QUANDT and DE HULTQUIST. Characterization of NADPH-dependent methemoglobin reductase as a heme-binding protein present in erythrocytes and liver. *Proc Natl Acad Sci USA* [online] . 1992, vol 89, no. 6, pp. 2130-4, also available from < <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC48610/?tool=pubmed> >. ISSN 0027-8424.
6. ↑ CORTAZZO, Jessica A and Adam D LICHTMAN. Methemoglobinemia: a review and recommendations for management. *J Cardiothorac Vasc Anesth* [online] . 2014, vol 28, no. 4, pp. 1043-7, also available from < <https://www.ncbi.nlm.nih.gov/pubmed/23953868> >. ISSN 1053-0770 (print), 1532-8422.
7. ↑ LEDVINA, M. Rapid spectrophotometric determination of carbonylhemoglobin in blood. *Biochem Clin Bohemoslov*. 1987, vol 16, pp. 493-495, ISSN 0139-9608.
8. ↑ Jump up to: a b ŠEBKOVÁ, Sylva. *Iron poisoning* [online]. © 2003. Last revision 2003-10-06, [cited. 2021-08-16]. < <http://medicina.cz/clanky/5819/34/Otrava-zelezem/> >.

References

- BUBNOVÁ, Eva, Alena BUDĚŠÍNSKÁ and Jana STRÍBRNÁ. *Practical exercises in medical chemistry and molecular biology*. 1st edition. Prague: Karolinum, 1998. 200 pp. ISBN 80-7184-725-9 .
- BURTIS, Carl A and Edward R ASHWOOD. *Tietz textbook of clinical chemistry*. 2nd edition. Philadelphia: Saunders, 1994. 2326 pp. ISBN 0-7216-4472-4 .
- ČERMÁK, J. Current possibilities of therapeutic influencing of conditions with iron deficiency or excess in the organism. *Remedia*. 1994, vol. 4, pp. 254-259, ISSN 0862-8947.
- DOLEŽALOVÁ, Věra, et al. *Laboratory technique in clinical biochemistry and toxicology*. 4th edition. Brno: Institute for Further Education of Healthcare Workers, 1995. 286 pp. ISBN 80-7013-198-5 .
- DZURIK, Rastislav. *Standard clinical biochemical diagnostics*. 2nd edition. Martin: Osveta, 1996. 464 pp. ISBN 80-217-0256-7 .
- CHROMY, Vratislav. *Analytical methods in clinical chemistry*. 1st edition. Brno: Masaryk University, Faculty of Science, 2000. 211 pp. ISBN 80-210-2363-5 .
- KAPLAN, Lawrence A, Amadeo J. PESCE and Steven C. KAZMIERCZAK. *Clinical chemistry*. 3rd edition. St. Louis: Mosby, 1996. 1064 pp. ISBN 0-8151-5243-4 .
- Team of authors. . *Medical chemistry and biochemistry*. 1st edition. Praha, Bratislava: Avicenum, Osveta, 1991. 237 s. 2. ISBN 80-201-0114-4 .
- KRAML, Jiri. *Instructions for practical exercises in medical chemistry and biochemistry*. 4th edition. Prague: Karolinum, 1991. 311 pp. ISBN 80-7066-453-3 .
- MASOPUST, Jaroslav. *Clinical biochemistry: requirements and evaluation of biochemical examinations*. 1st edition. Prague: Karolinum, 1998 .. 832 pp. ISBN 80-7184-650-3 .
- MEŠKO, Dušan. *Academy of Clinical Biochemistry*. 1st edition. Martin: Osveta, 1998. 1647 pp. ISBN 80-8063-005-4 .
- MURRAY, Robert K, et al. *Harper's biochemistry*. 2nd edition. Prague: H & H, 1998. 872 pp. ISBN 80-85787-

- RACEK, Jaroslav, et al. *Clinical biochemistry*. 1st edition. Prague: Galén, 1999. ISBN 80-7262-023-1 .
- SCHNEIDERKA, Petr. *Determination of analytes in clinical biochemistry*. 1st edition. Prague: Karolinum, 1999. 153 pp. Vol. 1. ISBN 80-7184-761-5 .
- SCHNEIDERKA, Petr. *Selected chapters from clinical biochemistry*. 1st edition. Prague: Karolinum, 1998. 119 pp. ISBN 80-7184-505-1 .
- ŠTÍPEK, Stanislav. *Brief toxicology*. 1st edition. Prague: Medprint, 1997. 87 pp. ISBN 80-902036-4-7 .
- TÁBORSKÁ, Eva and Josef TOMANDL, et al. *Biochemistry*. 1st edition. Brno: Masaryk University, 1998. 160 pp. ISBN 80-210-1736-8 .
- TÁBORSKÝ, Otto, et al. *Methods of clinical biochemistry: intended for students of the Faculty of Science*. 2nd edition. Prague: SPN, 1990. 158 p.
- THOMAS, Lothar, et al. *Clinical Laboratory Diagnostics*. 1st edition. Frankfurt am Main: TH-Books Verlagsgesellschaft, 1998. 1727 pp. ISBN 3-9805215-4-0 .
- ZIMA, Tomas, et al. *Laboratory diagnostics*. 1st edition. Prague: Galén, 2003. 728 pp. ISBN 80-7262-201-3 .