

Hypothalamo-pituitary system

The **hypothalamo-pituitary system** is part of the neuroendocrine system (the system connecting nerve signals to endocrine signals) and controls the other endocrine glands. It consists of two parts: the hypothalamus and the pituitary gland.

Hypothalamus

 For more information see *Hypothalamus*.

The **Hypothalamus** is a part of the diencephalon containing the control center of the body's autonomic nervous system. It creates hormones by itself, namely:

- **liberins** and **statins**, which regulate the release of hormones from the adenohypophysis;
- **antidiuretic hormone** (from nucleus supraopticus hypothalami) and **oxytocin** (from nucleus paraventricularis hypothalami).

It is associated with the pituitary gland, namely:

- **vascular** – it is connected to the adenohypophysis by a portal system. **A. hypophysialis superior**, a branch of a. carotis interna, breaks up into the primary capillary plexus at the level of eminentia mediana hypothalami, where liberins and statins are released into it. The primary capillary plexus rejoins into the portal veins, which subsequently disintegrate into the secondary plexus. This is located around the adenohypophysis, where liberins and statins are absorbed. On the contrary, hormones produced by the adenohypophysis itself pass into the blood. Vv leave from the secondary braid. hypophysiales to sinus cavernosus. This phenomenon, when the capillary plexus is inserted between vessels of the same type, is called **rete mirabile**, in this case it is **rete mirabile venosum** (the plexus is inserted between the veins).
- **nervously** – it is connected to the neurohypophysis by unmyelinated axons of neurons of the hypothalamic nuclei (nucleus supraopticus et paraventricularis). These axons form the stalk of the pituitary gland and are used to transport antidiuretic hormone ADH, vasopressin) and oxytocin. to the posterior lobe of the pituitary gland. The pathway formed by these axons is called the **hypothalamo-hypophysialis tract**.



Magnetic resonance imaging of the brain; the arrow points to the hypothalamus




Diagram of the hypothalamo-pituitary system

Hypophysis


 For more information see *Hypophysis*.

The **pituitary gland** is an oval extension of the end of the infundibular process of the hypothalamus. It is located in the sella turcica of the sphenoid bone and is composed of two anatomically, functionally and developmentally different parts – the **adenohypophysis** (with an epithelial structure) and the **neurohypophysis** (with a neural structure). The neurohypophysis and the adenohypophysis are connected to each other by a short portal circulation (*a. hypophyseal inferior*).

Adenohypophysis

The adenohypophysis is the anterior lobe of the pituitary gland. Hormone production by the adenohypophysis is regulated by liberins and statins from the hypothalamus. It is vascularly connected to it by the hypophyseal portal circulation. It consists of three parts

- **pars distalis** – the main part of the adenohypophysis responsible for the production of most hormones, formed by beams of chromophobe and chromophile cells. Chromophilic cells are further divided into **acidophilic** and **basophilic** cells.
- **pars intermedia** – poorly developed part formed mainly by basophilic cells.
- **pars tuberalis** – mainly contains chromophobe and Rathke cyst cells



1 – chiasma opticum, 2 – 3 – pars tuberalis, 4 – pars distalis, 5 – 6 – pars nervosa, 7a – 7b – 8 – 9 – 10 – neurosecretory neurons, 11 – nucleus paraventricularis, 12 – nucleus supraopticus, 13 – pars intermedia

Hormones of the adenohypophysis

Acidophilic cells produce simple proteins and predominate in squamous epithelium, they include two main cell types:

- **somatotropic cells** – **somatotropin** (STH), growth hormone - supports proteosynthesis, and thus individual growth, acts on the proliferation of chondrocytes in the growth plates;
- **mammotropic cells** – **prolactin** (PRL), stimulates mammary gland development and milk production.

Basophilic cells of the pars distalis secrete glycoproteins – they are PAS positive:

- **thyrotropic cells** – **thyrotropic hormone** (TSH), stimulates the synthesis and release of thyroid hormones;
- **gonadotropic cells** – **follicle-stimulating hormone** (FSH), in women it stimulates ovarian follicle maturation, in men it supports spermatogenesis and **luteinizing hormone** (LH), inhibits egg growth and increases the effect of estradiol on the storage and consumption of cholesterol in granulosa lutein cells, supports the production of sex hormones;
- **corticotropic cells** – **adrenocorticotrophic hormone** (ACTH), stimulates the growth of the adrenal cortex and the production of glucocorticoids.

In the basophilic cells of the pars intermedia, it is additionally formed:

- **melanocyte-stimulating hormone** (MSH) – stimulates melanocytes to produce melanin and transfer its granules to keratinocytes in the skin and hair.

Neurohypophysis

The neurohypophysis, or posterior lobe of the pituitary gland, is made up of axons of neurosecretory neurons and glial cells (so-called pituicity). It is nervously connected to the hypothalamus and through this route it also receives antidiuretic hormone and oxytocin, which are only stored in the pituitary gland and released into the blood by exocytosis after stimulation by an action potential.

- oxytocin – induces contractions of the uterus during childbirth and contraction of the milk ducts during breastfeeding.
- antidiuretic hormone – increases reabsorption of Na^+ and water in the distal tubule and collecting duct of the kidneys.

Links

Related articles

- Diseases of the hypothalamic-pituitary system
- Hypothalamus
- Hypophysis

References

- KITTNAR, Otomar. *Lékařská fyziologie*. 1. edition. Praha : Grada, 2011. 790 pp. ISBN 978-80-247-3068-4.
- ČIHÁK, Radomír – GRIM, Miloš. *Anatomie*. 2. upr. a dopl edition. Praha : Grada Publishing, 2002. 470 pp. vol. 2. ISBN 80-247-0143-X.
- KONRÁDOVÁ, Václava – UHLÍK, Jiří – VAJNER, Luděk. *Funkční histologie*. 2. edition. Jinočany : H & H, 2000. 291 pp. ISBN 80-86022-80-3.
- JUNQUEIRA, L. Carlos – CARNEIRO, Chosé. *Základy histologie*. 7. edition. Jinočany : H&H, 1999. ISBN 8085787377.