


CHA2DS2 - VASc score



CHA₂DS₂-VASc score is a scale used to 'assess the risk of thromboembolic complications' in patients with 'atrial fibrillation'. It replaced the older version of the 'CHADS₂' score, which underestimated some indications of anticoagulants compared to this version. In secondary prevention of stroke, the patient automatically has at least 2 points (risk of recurrence > 10% / year) and is indicated for anticoagulant therapy.

C	cardial failure selhání (EF < 40%)	1
H	hypertension	1
A₂	age ≥ 75	2
D	diabetes mellitus	1
S₂	stroke	2
V	vascular diseases (heart attack, ischemic disease of the lower limbs..)	1
A	age 65–74	1
Sc	sex category	1



Embolism in atrial fibrillation

- The maximum number obtained is '9', when 'the risk of thromboembolism is' highest.
- Patients with a 'score ≥ 2' are indicated for p.o. anticoagulant treatment.
- In patients 'at risk 1', anticoagulant therapy is rather recommended.
- Patients 'without risk factors' do not require treatment or are given antiplatelet agents (ASA).

Calculator

MediaWiki:Lékařská kalkulačka/Antikoagulace

Links

Related articles

- Atrial fibrillation
- Embolization
- Anticoagulants

Sources

- CHA₂DS₂-VASc score (stroke manual) (<http://www.cmp-manual.cz/914-CHADS2-score.html>)
- new anticoagulants in the prevention of TE complications of FS (<http://www.practicus.eu>)

External links

- Chadsvasc.org (<https://www.chadsvasc.org/>)