

# Classification of otitis media

We distinguish between inflammations:

- **purulent and non-purulent ;**
- **acute and chronic .**Acute otitis media-exudate arches the membrane laterally

The basic 4 types of mesotitids are:

1. *acute tubotympanic catarrh* (catarrhus tubae auditivae acutus, hydrotympanum);
2. *chronic tubotympanic catarrh* (catarrhus tubotympanalis chronicus, secretory mesotitis, otitis media secretorica - OMS, glue ear);
3. *acute purulent mesotitis* (otitis media suppurativa acute, otitis media acute - OMA)
4. *chronic purulent mesotitis* (otitis media suppurativa chronica, otitis media chronica)



Acute otitis media-exudate arches the membrane laterally

## Links

### Related Articles

- Complications of otitis media
- Otitis media (guide)

### References

- KLOZAR, Jan, et al. *Speciální otorinolaryngologie*. 1. edition. Praha : Galén, 2005. 224 pp. ISBN 80-7262-346-X.
- CHROBOK, Viktor – MICHÁLEK, Roman. Diagnostika a léčba středoušního zánětu z pohledu ORL lékaře. *Lékařské listy* [online]. 2009, vol. 15, p. 30, Available from <<https://zdravi.euro.cz/clanek/priloha-lekarske-listy/diagnostika-a-lecba-stredousniho-zanetu-z-pohledu-orl-lekare-445464>>. ISSN 1214-7664.