

# Combination of NSAID

## Input

Prescribe a combination of NSAID (analgesic – antipyretic, Non-steroidal anti-inflammatory drugs) and substances enhancing the analgesic effect for an adult patient in divided powders (magistraliter).

In practice, the combination is quite often prescribed magistraliter, especially in older patients, in whom it is necessary to give lower doses, which are not produced as HVLP.

## Solution

### Recipe

Paracetamol	0.4
Codeine dihydrogenphosphatis	0.01
Caffeine	0.03

Bowl fiat pulvis.

Da tales doses No X (decem) ad capsulas gelatinosas.

Signa *1 tbl. in the morning, 1 tsp. at noon, 1tbl. evening.*

Rp.

**Acidi acetylsalicylici** 0.48  
Codeini phosphatis sesquihydrici 0.02  
M. f. pulv.  
D. t. d No. XX (vigints) ad caps.  
S.: For difficulties, 1 capsule 3-4 times a day.

Rp.

**Acidi acetylsalicylici** 9.60  
Codeini phosphatis sesquihydrici 0.40  
M. f. pulv.  
Wonder. in doses aequales No. XX (vigints) ad caps.  
S.: For difficulties, 1 capsule 3-4 times a day.

## Links

### Related Articles

- Ointment with 5% boric acid
- Atropine eye drops
- Opioid analgesic drops
- Keratolytic Ointment
- Paracetamol in suppositories
- Salinic laxans
- Eye drops with 1% pilocarpine
- Keratoplasty Ointment
- Liquid backfill with 5% tannin

### References

- VONDRÁČKOVÁ, Hana. *Master's recipe* [lecture for subject Pharmacology, specialization General medicine, 1. LF UK]. Praha. 2010-04-18.

### Recommended reading

- HEJLEK, Jan – JANÁT, Vratislav – JINDRA, Lumír, et al. *Prescriptiones magistrales*. 4th, edited edition. Avicenum, 1976. 442 pp.
- JANŮ, Michael – KOROUSOVÁ, Marie – PROCHÁZKOVÁ, Jindra, et al. *Individually prepared medicinal products*. 1. edition. Praha : Galén, 2009. 111 pp. Special purpose publication of the VFN hospital pharmacy in

