

# Cortico-dependent asthma

**Cortico-dependent asthma** is asthma in patients with severe forms of asthma (OIA), when systemic oral administration of corticoids is necessary.

About 5% of asthma patients fall into the difficult-to-treat asthma (OIA) category. Asthma cannot be achieved or permanently controlled in them.

Aggravation caused by reflux irritation , but also **reduced compliance and adherence** to treatment in severe **psychiatric** diseases can lead to OIA.

## Links

### related articles

Passwords:

- RABA
- SHEBA
- PAW

On the VPL portal:

- Bronchial asthma therapy
- Therapy of an attack of asthma bronchiale

In Wikiscripts:

- Astma
- Exercise: Asthma bronchiale
- Status asthmaticus
- Bronchial asthma therapy

### External links

[www.svl.cz/....astma-2008.pdf](http://www.svl.cz/....astma-2008.pdf) ([https://www.svl.cz/Files/nastenka/page\\_4771/Version1/astma-2008.pdf](https://www.svl.cz/Files/nastenka/page_4771/Version1/astma-2008.pdf))

### References

- SALAJKA, František. *Asthma bronchiale : Recommended diagnostic and treatment procedure for general practitioners* [online] . 1st edition. Prague: Society of General Medicine ČLS JEP, 2008. Also available from < <https://www.svl.cz/default.aspx/cz/spol/svl/default/menu/doporucenepostu/doporucenepostu5> >. ISBN 978-80-86998-26-8

Kategorie:Všeobecné praktické lékařství Kategorie:Pneumologie