

Dead fetus syndrome

When a dead fetus is detected during pregnancy or when it dies during childbirth, we speak of **dead fetus syndrome**. This threatens the mother's health and life. These are all deaths during pregnancy and childbirth (intrapartum death), when the weight of the fetus is greater than 500 g and none of the signs of life are present (heart beat, breathing movements, active muscle movements, screams).

Causes of intrauterine fetal death

The most common causes of antenatal fetal death ^[1]	Fetal causes	Placental and umbilical cord causes
Maternal causes	Fetal causes	Placental and umbilical cord causes
Preeclampsie, Eclampsia, HELLP syndrome	structural malformation	placental insufficiency with IUGR
diabetes mellitus	chromosomal aberrations	placenta praevia, vasa praevia
nephropathy, hepatopathy	congenital disorders of metabolism	placental abruption
cardiovascular and cardiopulmonary diseases	intrauterine infection	strangulation, prolapse of the umbilical cord
antiphospholipid syndrome	feto-fetal transfusion of twins	chorioamnitis
shock, sepsis, profusional bleeding	hydrops	hemangiomas, teratomas of the placenta; myxomas, hemangiomas of the umbilical cord

Diagnostics

Subjective feelings of a woman

- reduced frequency or absence of perception of fetal movements;
- vaginal bleeding or discharge;
- subfebrile;
- fatigue;
- breast tension laxity.

Objective diagnosis

Intrauterine fetal death is diagnosed based on the lack of fetal heart activity. Heart sounds are not heard during examination with a stethoscope, during Doppler ultrasonography or during cardiotocography (CTG). **If the dead fetus is present in the uterus for more than 7 days, a roof-like overlapping of the skull bones (Spalding's sign)** can be seen on the ultrasound image.

Risks of stillbirth syndrome

- **Disseminated intravascular coagulation** (DIC) – dysregulation of coagulation in the mother's bloodstream occurs after activation of the plasma coagulation system. The trigger is tissue factor, which is washed out on the basis of increased intra-amniotic pressure (on the basis of autolytic processes of the fetus) and a broken uteroplacental barrier.
- **Infection and sepsis** – at first intraovular infection becomes systemic, in the most severe cases septic shock and coagulation disorders occur. Antibiotics are administered prophylactically.
- **Maternal illness that led to the death of the fetus** – most often it is severe preeclampsia, HELLP syndrome, hepatorenal failure, cardiovascular or cardiopulmonary failure.

Therapeutic procedure

1. Inform the mother about the death of the fetus.
2. Immediate hospitalization at an obstetrics and gynecology clinic or, in case of expected complications, at a perinatology center.
3. Laboratory examination.
 For more information see *Routine lab tests in a woman with a dead fetus*.
4. Prophylaxis:
 - prevent the occurrence of DIC by applying low-molecular-weight heparin (LMWH) according to body weight 0.2-0.4 ml/24 h, further according to the control of hemocoagulation parameters ;
 - administration of broad-spectrum ATBs as infection prevention.
5. Induction of labor - we choose between classic (disruption of the sac of membranes) and prostaglandin induction;

- classic induction – induction associated with disruption of the sac of membranes is chosen when the woman is bleeding, when the placenta is separated prematurely, the condition is a permeable cervix; after disruption of the sac, we can continue with the infusion of oxytocin;
- prostaglandin induction – prostaglandin E₂ can be applied locally in the form of a gel (*Prepidil*) or as tablets (*Prostin*).

Management of childbirth

We primarily deliver vaginally. Stillbirth induction is usually quicker and easier. More caution and observation of the patient during the entire delivery is always appropriate. After the start of regular uterine activity, we apply epidural analgesia and spasmolytics, or smaller doses of opiates. We do not perform an episiotomy, especially if the cranial vault is collapsed. Delivery of the fetus by caesarean section is indicated in the event of an urgent situation (e.g. heavy bleeding of the mother after placental abruption). We deliver the fetus and placenta as one unit without cutting the umbilical cord.

Links

related articles

- Preeclampsia
- Eclampsia
- HELLP syndrome

Reference

1. ČECH, Evžen. *Porodnictví*. 2. edition. Praha : Grada Publishing, Praha : Avicenum, 2006. pp. 355-357. ISBN 8024713039.

External links

- JEŽOVÁ, Marta – HOTÁRKOVÁ, Sylva – MŮČKOVÁ, Katarína, et al. *Hypertextový atlas fetální patologie : Multimediální podpora výuky klinických a zdravotnických oborů* [online]. Portál Lékařské fakulty Masarykovy univerzity [online], ©2008. The last revision 2.2.2010, [cit. 26.11.2011]. <<http://portal.med.muni.cz/clanek-463-hypertextovy-atlas-fetalni-patologie.html>>.
- JEŽOVÁ, Marta – HOTÁRKOVÁ, Sylva – MŮČKOVÁ, Katarína, et al. *Hypertextový atlas novorozenecké patologie : Multimediální podpora výuky klinických a zdravotnických oborů* [online]. Portál Lékařské fakulty Masarykovy univerzity [online], ©2010. The last revision 27.9.2011, [cit. 26.11.2011]. <<http://portal.med.muni.cz/clanek-527-hypertextovy-atlas-novorozenecke-patologie.html>>.

References

- ČECH, Evžen. *Porodnictví*. 2. edition. Praha : Grada Publishing, Praha : Avicenum, 2006. pp. 355-357. ISBN 8024713039.