

Oxidative phase of the pentose cycle

Under construction / Forgotten

This article was marked by its author as *Under construction*, but the last edit is older than 30 days. If you want to edit this page, please try to contact its author first (you will find him in the history (https://www.wikilectures.eu/index.php?title=Oxidative_phase_of_the_pentose_cycle&action=history)). Watch the page as well. If the author will not continue in work, remove the template `{{Under construction}}` and the page.

Last update: Wednesday, 15 Feb 2023 at 12.54 am.

`{{editorcheck}}`

In addition to AcCoA, **propionyl-CoA** is also formed by oxidation of odd-chain fatty acids. This is first **carboxylated to methylmalonyl-CoA**, which is converted to **succinyl-CoA** - an intermediate of the **Krebs cycle**. Through conversion to oxaloacetate, it can participate in **gluconeogenesis** - glucose can be synthesized from these fatty acids. However, very few fatty acids with an odd number of carbon atoms are found in the body.

Links

Related Articles

- Oxidation of very long chain fatty acids
- Oxidation of unsaturated fatty acids
- Regulation of beta-oxidation of fatty acids
- Lipid breakdown and ketone body metabolism

Source

Fontana J., Trnka J., Maďa P., Ivák P. et al.: Transformation of substances and energy in the cell. In: Functions of cells and the human body : Multimedia scripts. Available online from: [1] (<http://fbt.cz/skripta/ii-premena-latek-a-energie-v-bunce/>)


The **pentose cycle** is a catabolic event that provides reduced cofactors NADPH and five-carbon saccharides, or pentoses. It is a metabolic conversion of glucose, the goal of which is not the creation of ATP.

Course of the oxidative phase of the pentose cycle

During the *oxidative phase* of the pentose cycle, the glucose-6-P molecule is oxidized to the ribulose-5-P molecule. At the same time, CO₂ is released and two NADPH + H⁺ molecules are obtained.

File:Oxidation.jpg
Scheme of the oxidative phase of the pentose cycle

Its course is summarized by the following equations:



Of the reactions of the first phase, the initial reaction catalyzed by `glucose-6-phosphate dehydrogenase` is important. This **irreversible reaction is the main regulatory step of the pentose cycle**.

The rate of the oxidative phase of the pentose cycle

The **speed** of the entire metabolic pathway depends on the activity of two dehydrogenation reactions, which depend on the availability of NADP⁺ (i.e. the oxidized form of the coenzyme). With a lack of NADP⁺, the rate of the pentose cycle decreases, in other words: an excess of NADPH "slows down" the oxidative phase of the pentose cycle.

Links

Related Articles

- Pentose cycle
- NADPH

External links

- Pentose cycle (Czech Wikipedia) (https://cs.wikipedia.org/wiki/Pent%C3%B3zofosf%C3%A1tov%C3%BD_cyklus%7C)